



Icelandic tephrochronology - matching the provenance of proximal and distal volcanic glasses using La-ICPMS trace element data

Martin Menzies, E. Tomlinson, Wolfgang Muller, T. Thordarson, Christine Lane, Vicky Smith, S. Blockley

► To cite this version:

Martin Menzies, E. Tomlinson, Wolfgang Muller, T. Thordarson, Christine Lane, et al.. Icelandic tephrochronology - matching the provenance of proximal and distal volcanic glasses using La-ICPMS trace element data. Iceland in the Central Northern Atlantic: hotspot, sea currents and climate change, May 2010, Plouzané, France. hal-00482055

HAL Id: hal-00482055

<https://hal.univ-brest.fr/hal-00482055>

Submitted on 7 May 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ICELANDIC TEPHROCHRONOLOGY - MATCHING THE PROVENANCE OF PROXIMAL AND DISTAL VOLCANIC GLASSES USING LA-ICPMS TRACE ELEMENT DATA

Martin Menzies¹, Emma Tomlinson¹, Wolfgang Muller¹, Thor Thordarson², Christine Lane³, Vicky Smith³, Simon Blockley⁴ and the RESET Consortium⁵

¹Earth Sciences, Royal Holloway University of London, Egham

²School of GeoSciences, University of Edinburgh, Edinburgh

³Research Laboratory for Archaeology & the History of Art, Oxford

⁴Department of Geography, Royal Holloway University of London, Egham

⁵RESET WWW site : <http://c14.arch.ox.ac.uk/reset/>

Abstract

Thorarinsson (1944) pioneered the use of tephrochronology and its application in NW Europe. The basis of this chronological tool is the use of time-parallel marker tephra (i.e., ash <2mm) horizons to correlate proximal volcanic deposits with distal tephtras found in marine-lacustrine-ice cores and terrestrial sequences. The trace element geochemistry of volcanic glasses [determined by LA-ICPMS] offers an improved diagnostic tool for matching the juvenile components of proximal and distal tephtras and has considerable potential for assigning provenance to cryptotephtras (<100 microns) provided they are of sufficient size and thickness.

Introduction

In the context of the last 100ka western Europe offers a unique opportunity to use recurrent explosive volcanic activity and widespread tephra dispersal as a basis for evaluating the temporal and spatial relationships between the migration of ancient modern humans (AMH) and abrupt environmental transitions (AETs). Tephtras in (a) archaeological sites hold the key to AMH migration patterns and (b) lacustrine, marine and ice core constrain AETs. Juvenile (magmatic glass) compositions are forthcoming from (a) pumice or scoria clasts in pyroclastic fall deposits (b) pumice or fiamme in proximal pyroclastic flow deposits, (c) glass adhering to entrained xenoliths, and (d) cryptotephtras at distal sites (i.e., < 100 microns). EMPA & LA-ICPMS techniques (e.g., Tomlinson et al 2010a) are applied to all proximal and distal juvenile glass with SIMS utilized where particles are too small/thin. The provenance of these distal tephtras is paramount and is defined by matching the geochemistry of distal juvenile clasts with the RESET geochemical database for proximal European tephtras. Once diagnostic chemistries are matched, ⁴⁰Ar/³⁹Ar ages (K-rich phenocrysts) from proximal sites link the continental-lacustrine-marine archives. This time lattice is refined with the addition of temporal detail from varve counting (lacustrine), ¹⁴C, U-Th dating (archaeological sites), & biostratigraphy (marine).

Katla & the Vedde ash (ca. 11ka) : proximal-distal correlation

The Vedde Ash has been reported from distal terrestrial, marine and ice cores across Northern Europe and forms an important stratigraphic marker for palaeoclimatic reconstructions across the North Atlantic (e.g., Mangerud et al. 1984; Lowe and Turney, 1997; Wastegård et al., 2000; Davies et al., 2005). The Vedde ash has been dated close to its type site in Western Norway, giving a ¹⁴C age of 10,310 ± 50 ¹⁴C years or 11,841 to 12,383 cal years BP (Birks et al., 1996). The Vedde Ash is a bimodal rhyolitic (clear shards) and basaltic (brown shards) tephra layer believed to have been erupted from the Katla volcano in Southern Iceland (Lacasse et al., 1995). However, distal tephra correlations are solely based on somewhat equivocal data including glass shard morphology and major element EMPA chemistry indicating a basalt-rhyolite magmatic lineage. Lacasse et al (1995) identified the Sólheimar ignimbrite, erupted from Katla, as the likely source of the Ash Zone 1 tephra again on the basis of total alkali-silica (TAS) classification and feldspar compositions. These may not be diagnostic in the context of Icelandic magmatism and the potential exists for confusion between distal basalt-rhyolite cryptotephra and the many basalt-rhyolite centres on Iceland that have erupted over the last 100ka. Tomlinson et al (2010b) undertook trace (LA-ICP-MS) element analysis of volcanic glasses from a proximal volcano-stratigraphic section of the Sólheimar ignimbrite and distal glasses from the type Vedde site at Ålesund in Western Norway (Mangerud et al., 1984). The rhyolitic (SiO₂>68%) Sólheimar pumice and obsidian and the distal Vedde clear glass shards have restricted and



overlapping trace element compositions that demonstrate a close correlation between the Sólheimir ignimbrite and the Vedde Ash. Importantly Tomlinson et al (2010b) highlighted key differences between the Sólheimir rhyolite and other rhyolites from the Icelandic Southern Transition Zone such that provenance can be assigned with greater confidence. *They conclude that the proximal rhyolitic component of the Sólheimir ignimbrite is compositionally identical to the distal rhyolitic clasts from the classic Vedde fall locality in terms of both major and trace element geochemistry.* Magmatic variability exists along 1500 km of the tephra dispersal axis between Katla and Vedde with greater magmatic representation in the distal Vedde tephra (i.e. fall deposit) than at the proximal "bimodal" site at Sólheimir which is dominantly basaltic-andesite fall and rhyolitic flows deposits. Similar along-axis variability over 3000km was reported from continental-marine correlations between Afro-Arabia and the Indian Ocean (Ukstins-Peate et al 2003, 2008). Along axis differences appear to be characterised by (a) the most evolved magmas being located at distal sites, and (b) bimodality being restricted to proximal sites.

Thorsmork & Ash Zone 2 (ca. 57ka) – continental-marine correlation

Ash zone 2, an alkali rhyolite tephra layer found in north Atlantic and Arctic marine cores (Lacasse & Garbe-Schoonberg 2001) and the GISP2 Greenland ice core (Ram et al 1996), has been correlated with the Thorsmork ignimbrite. This provenance match is based on a combination of major element data and tephra stratigraphy. Tindfjallajökull, the source of the Thorsmork ignimbrite, is a bimodal basalt-rhyolite volcano in the Eastern Volcanic Zone Iceland, represented by a series of pyroclastic flow deposits. The homogenous rhyolite pumice dominates the deposit, which also contains subordinate basalt scoria (Jørgensen 1980). These two compositional components are also represented in the fiamme but there is no textural evidence for mixing or mingling between the basalt and rhyolite components. The Thorsmork ignimbrite has been the focus of $^{40}\text{Ar}/^{39}\text{Ar}$ dating (Sigurdsson et al 1998, Storey and Stecher 2009) with a robust age of 57ka (W. McIntosh 2010 pers comm). Tomlinson et al (2010a) report LA-ICP-MS and AES data for the Thorsmork rhyolite and its juvenile component. They argue that the bimodal Thorsmork ignimbrite is the product of physical mixing of basaltic and rhyolitic magmas. Furthermore Tomlinson et al (2010a) compare other potential volcanic sources from the Icelandic Southern Transition Zone (Hekla, Torfajökull, Oaerfajökull, Tindfjallajökull). Rhyolite glass from the Thorsmork Ignimbrite can be distinguished from these other Icelandic rhyolites, because the major differences between the Icelandic rhyolites are driven by high-level processes in the magma chamber, principally phenocryst fractionation. Tomlinson et al (2010a) compare the Thorsmork glass chemistry with published trace element data for ash zone 2 tephra (SU9029) from the Irminger Basin, North Atlantic (Lacasse & Garbe-Schoonberg 2001). They conclude that *ash zone 2 is indistinguishable from the proximal Thorsmork juvenile clasts and that Ash Zone 2 is likely to be a distal sample of the Thorsmork Ignimbrite.*

Summary

Volcanic glass geochemistry offers a robust basis for matching proximal juvenile clasts, from primary fall & flow deposits, with distal tephras and cryptotephras [$<100\mu\text{m}$]. Glass chemistries from two Icelandic volcanoes have been used to demonstrate : (a) proximal-distal correlations over 1500km with Katla confirmed as the source of the "Vedde Ash" in Norway (11ka), and (b) continental-marine correlations with Thorsmork (57ka Tindfjallajökull) shown to be the source of Ash Zone 2 found in the North Atlantic marine cores.

Acknowledgements

The RESET Consortium (2008-2012) is funded by the NERC (UK) and the CNRS supported attendance at the Ecole Thématique CNRS Summer School "Iceland and the Northern Atlantic" in Plouzané France 2010.



References

- Birks, H.H., Gulliksen, S., Hafliðason, H., Mangerud, J., and Possnert, G., 1996, New radiocarbon dates for the Vedde ash and the Saksunarvatn ash from western Norway: *Quaternary Research*, **45**, 119-127.
- Davies, S.M., Hoek, W.Z., Bohncke, S.J.P., Lowe, J.J., O'Donnell, S.P., and Turney, C.S.M., 2005, Detection of Lateglacial distal tephra layers in the Netherlands: *Boreas*, **34**, 123-135.
- Jørgensen, K., 1980. The Thorsmork Ignimbrite - an unusual comenditic pyroclastic flow in southern Iceland. *Journal of Volcanology and Geothermal Research*, **8**, 7-22.
- Lacasse, C. and Garbe-Schoonberg, C.-D., 2001. Explosive silicic volcanism in Iceland and the Jan Mayen area during the last 6 Ma: sources and timing of major eruptions. *Journal of Volcanology and Geothermal Research*, **107**, 113-147
- Lacasse, C., Sigurdsson, H., Johannesson, H., Paterne, M., and Carey, S., 1995, Source of Ash-Zone-1 in the North-Atlantic: *Bulletin of Volcanology*, **57**, 18-32
- Lowe, J.J., and Turney, C.S.M., 1997, Vedde ash layer discovered in a small lake basin on the Scottish mainland: *Journal of the Geological Society*, v. **154**, 605-612.
- Mangerud, J., Lie, S.E., Furnes, H., Kristiansen, I.L., and Lømo, L., 1984, A Younger Dryas Ash Bed in western Norway, and its possible correlations with tephra in cores from the Norwegian Sea and the North Atlantic: *Quaternary Research*, **21**, 85-104.
- Ram M., Donarummo J. and Sheridan M., 1996. Volcanic ash from Icelandic similar to 57,300 yr BP eruption found in GISP2 (Greenland) ice core. *Geophysical Research Letters*, **23**, 3167-3169
- Sigurdsson, H., McIntosh, W.C., Dunbar, N., and Lacasse, S.N. (1998) Thorsmork Ignimbrite in Iceland: possible source of North Atlantic ash. *EOS Transactions AGU*.
- Storey, M. and Stecher, O. (2009) 40Ar/39Ar Age of the Thorsmörk ignimbrite, Iceland and correlation with North Atlantic Z2 ash: Cross-calibration of ~60,000 counted annual layers in the Greenland ice cores. *EOS Transactions AGU*.
- Thorarinsson, S. (1944) Tefrokronologiska studier på Island. *Geografiska Annaler*, **26**, 1-217.
- Tomlinson, E., Thordarson, T., Lane, C., Blockley, S., Muller, W., and Menzies, M.A. (2010a) Micro-analysis of tephra by LA-ICP-MS. *Chemical Geology* (in review)
- Tomlinson, E., Thordarson, T., Lane, C., Blockley, S., Muller, W., and Menzies, M.A. (2010b) The Solheimer ignimbrite (Katla, southern Iceland) and the source of the "Vedde Ash" stratigraphic marker tephra. *Geology* (in preparation)
- Ukstins Peate, I. et al., 2003. Correlation of Indian Ocean tephra to individual Oligocene silicic eruptions from Afro-Arabian flood volcanism. *Earth and Planetary Science Letters*, **211**, 311-327.
- Ukstins Peate, I., Kent, A.J.R., Baker, J.A. and Menzies, M.A., 2008. Extreme geochemical heterogeneity in Afro-Arabian Oligocene tephra: Preserving fractional crystallization and mafic recharge processes in silicic magma chambers. *Lithos*, **102**, 260-278.
- Wastegård, S., Wohlfarth, B., Subetto, D.A., and Sapelko, T.V., 2000, Extending the known distribution of the Younger Dryas Vedde Ash into northwestern Russia: *Journal of Quaternary Science*, **15**, 581-586.



RESET Response of Humans to Abrupt Environmental Transitions